

Non-local current correlations in ferromagnet/superconductor nanojunctions

C. J. Lambert^a

Department of Physics, Lancaster University, Lancaster, LA1 4YB, UK

J. Koltai, J. Cserti^b

*Department of Physics of Complex Systems, Eötvös University
H-1117 Budapest, Hungary*

When two fully polarized ferromagnetic (F) wires with opposite polarizations make contact with a spin-singlet superconductor, a potential-induced current in wire 1 induces a non-local current of equal magnitude and sign in wire 2. The magnitude of this current has been studied in the tunneling limit and found to decay exponentially with the distance between the contact. In this paper we propose a new structure in which this novel non-local effect is increased by orders of magnitude. We study the spin-dependent electronic transport of a diffusive nanojunction and demonstrate that when a normal diffusive region is placed between the F leads and superconductor, the non-local initially increases with the separation between the F leads, achieving a maximum and decays as a power law with increasing separation.

Experimental studies of electronic transport properties of nanostructures containing both ferromagnets (F) and superconductors (S)^{1,2,3,4,5,6,7,8,9,10,11} reveal novel features, not present in normal-metal/superconductor (N/S) junctions, due to the suppression of electron-hole correlations in the ferromagnet. When spin-flip processes are absent, further effects are predicted, including the suppression of the conventional giant magnetoresistance ratio in diffusive magnetic multilayers¹² and the appearance of non-local currents when two fully polarized ferromagnetic (F) wires with opposite polarizations make contact with a spin-singlet superconductor^{13,14}. The latter effect has been highlighted, because of interest in the possibility of generating entangled pairs of electrons at a superconductor (S) interface^{15,16,17}. A recent study of such a junction in the tunneling limit predicts that the magnitude of the non-local current will decrease exponentially with the distance between the F contacts. It is therefore of interest to ask how the magnitude of this novel effect can be enhanced.

In this paper we propose a hybrid nanostructure in which the non-local current is enhanced by orders of magnitude compared with the geometry of ref.¹³. The proposed structure is sketched in figure 1 and comprises two clean F wires, each of width M_f , separated by a distance M , in contact with a diffusive, normal metallic region of area $A = L(2M_f + M)$, which in turn makes contact with a spin-singlet superconductor.

In the linear response limit, the current-voltage relation of a such a hybrid structure connected by non-superconducting wires to normal reservoirs was first presented in ref¹⁸. If I_1, I_2 are the currents leaving reservoirs 1 and 2, and v_1, v_2

their respective voltages, then at zero temperature¹⁸

$$I_1 = [2e^2/h][(N - R_0 + R_a)(v_1 - v) + (T'_a - T'_0)(v_2 - v)] \quad (1a)$$

and

$$I_2 = [2e^2/h][(T_a - T_0)(v_1 - v) + (N - R'_0 + R'_a)(v_2 - v)], \quad (1b)$$

where v is the condensate potential. In this expression R_0 is the coefficient for an electron from reservoir 1 to be reflected as an electron back into reservoir 1, while T_0 is the coefficient for an electron from reservoir 1 to be transmitted as an electron into reservoir 2. R_a is the coefficient for an electron from reservoir 1 to be Andreev reflected as a hole into wire 1 and T_a is the coefficient for an electron from reservoir 1 to be transmitted as a hole into wire 2. Finally R'_0 , T'_0 , R'_a , T'_a are the corresponding coefficients for electrons originating from reservoir 2. All coefficients are evaluated at the Fermi energy and are obtained by summing over the elements of sub-matrices of the quantum mechanical scattering matrix S . As a consequence of unitarity of S , they satisfy $N = R_0 + R_a + T_a + T_0 = R'_0 + R'_a + T'_a + T'_0$, where N is the number of open scattering channels in the wires.

As discussed in ref¹⁹, equations (1) are a convenient starting point for describing a wide variety of transport phenomena in hybrid superconducting nanostructures. For example if lead 2 is employed as a voltage probe, with I_2 set to zero, then equation (1b) yields

$$(v_2 - v)/(v_1 - v) = (N - R'_0 + R'_a)/(T_0 - T_a). \quad (2)$$

In the absence of Andreev processes, this ratio is positive. However if Andreev transmission dominates normal transmission, the ratio can be negative, leading to novel negative 4-probe conductances in hybrid superconducting structures²⁰. The appearance of non-local currents is similarly obtained from equations (1) by setting $v_2 - v = 0$, which yields

$$I_1 = [2e^2/h][N - R_0 + R_a](v_1 - v) = [2e^2/h][2R_a + T_0 + T_a](v_1 - v) \quad (3)$$

and

$$I_2 = [2e^2/h][T_a - T_0](v_1 - v). \quad (4)$$

When all Andreev processes are absent, this yields the expected result $I_2 = -I_1 = -(2e^2/h)T_0(v_1 - v)$, whereas when Andreev reflection and normal transmission are completely suppressed, $I_2 = +I_1 = (2e^2/h)T_a(v_1 - v)$. The latter occurs when the F wires are completely spin polarized and the polarization in wire 1 is the opposite of that in wire 2. More generally I_1 and I_2 will have opposite signs whenever Andreev transmission dominates normal transmission. As noted in ref²⁰, this can occur even if the current carrying wires are not ferromagnetic.

In the absence of a normal region in front of the superconductor (ie $L = 0$) and in the presence of tunnel junctions at the F-S interfaces, the analysis of ref¹³ predicts that both I_1 and I_2 decay as $\exp(-(2L/\pi\xi))$, where ξ is the superconducting coherence length. To analyze this structure for finite L we solve the Bogoliubov - de Gennes equation on a square tight binding lattice and compute the scattering matrix using an exact recursive Green's function technique.

We start by defining a tight-binding lattice of sites with the geometry of figure 1. Each site is labeled by a lattice vector $\vec{l}$ and possesses particle (hole) degrees of freedom $\psi^\sigma(\vec{l})$ ($\phi^\sigma(\vec{l})$), with spin $\sigma = \pm 1$. Ferromagnetism is incorporated via a Stoner model, with an exchange splitting $h_0(\vec{l})$ on site $\vec{l}$. In the presence of local s-wave pairing described by a superconducting order parameter $\Delta(\vec{l})$, the Bogoliubov equation takes the form

$$E\psi^\sigma(\vec{l}) = [\epsilon(\vec{l}) - \sigma h_0(\vec{l})]\psi^\sigma(\vec{l}) - \sum_{\vec{\delta}} \gamma \psi^\sigma(\vec{l} + \vec{\delta}) + \sigma \Delta(\vec{l}) \phi^{-\sigma}(\vec{l}) \quad (5)$$

$$E\phi^\sigma(\vec{l}) = -[\epsilon(\vec{l}) - \sigma h_0(\vec{l})]\phi^\sigma(\vec{l}) + \sum_{\vec{\delta}} \gamma \phi^\sigma(\vec{l} + \vec{\delta}) - \sigma \Delta^*(\vec{l}) \psi^{-\sigma}(\vec{l}), \quad (6)$$

where $\vec{l} + \vec{\delta}$ sums over the nearest neighbors of $\vec{l}$.

If $\vec{l}$ belongs to the normal diffusive region of figure 1, $\epsilon(\vec{l})$ is chosen to be a random number, uniformly distributed over the interval $\epsilon_0 - W/2$ to $\epsilon_0 + W/2$, whereas in the clean F and S regions $\epsilon_i = \epsilon_0$. In the S region, the order parameter is set to a constant, $\Delta(\vec{l}) = \Delta_0$, while in all other regions, $\Delta(\vec{l}) = 0$. If $\vec{l}$ is located in one of the clean ferromagnetic leads, then Stoner splitting is $h_0(\vec{l}) = h_j$, where $j = 1, 2$ labels the lead containing site $\vec{l}$. For aligned moments $h_1 = h_2 = \epsilon_M$ and for anti-aligned moments, $h_1 = -h_2 = \epsilon_M$. If $\vec{l}$ is not located in one of the ferromagnetic leads then $h_0(\vec{l}) = 0$. The nearest neighbor hopping element γ fixes the energy scale (ie the band-width), whereas ϵ_0 determines the band-filling. In what follows, to model the experimentally-relevant regime of $\Delta_0 \ll$ Fermi energy ($= 4\gamma - \epsilon_0$) and $M_f >$ Fermi wavelength, we set $\gamma = 1$, $\epsilon_0 = 0.2$, $\epsilon_M = 3.8$, $M_f = 10$ and $\Delta_0 = 0.1$.

By numerically solving the Bogoliubov - de Gennes equation for a given energy E , the scattering matrix S can be computed for a given realization of the disorder and average values of measurable quantities obtained ensemble averaging the quantity of interest. For a given value of disorder W , the elastic mean free path λ_{el} of the normal region is obtained by from a separate calculation of the ensemble-averaged dc conductivity of a rectangular normal bar, connected to normal reservoirs. λ_{el} is then obtained by comparing the result with the Drude formula. In what follows, the lattice constant is set to unity and W chosen such that $\lambda_{el} = 10$.

For $E = 0$, figures 2 and 3 show the dependence on L and M of the ensemble-averaged Andreev transmission coefficient in the presence of non-magnetic wires, while figures 4 and 5 show the corresponding quantities in the presence of fully spin-polarized wires with opposite magnetizations. For small L (eg $L = 1$) and finite M , figures 3 and 5 show that T_a is exponentially small, as predicted by ¹³. However as L is increased to a finite value, T_a shows a dramatic enhancement.

Although this structure is difficult to describe analytically, the qualitative behaviour of T_a is clear and can be understood in terms of a simple resistor model, in which T_a is proportional to the conductance of electrons from lead 1 travelling first to the N-S interface and then to lead 2. Consider for example figure 2, which for $M \neq 0$ shows the expected exponential suppression of T_a at $L = 0$. For small L , T_a increases linearly with L . The slope of this linear regime is independent of M for small $M < \lambda_{el}$, reflecting the ballistic nature of the N region in this limit, whereas

for the opposite limit of $M > \lambda_{el}$, the slope decreases with increasing M , reflecting the diffusive behaviour of the N region. In the diffusive regime, for $M \ll L$, this leads us to expect the T_a is proportional to $L/(M + 2M_f)$, whereas for $L \ll M$ it is proportional to $(M + 2M_f)/L$, which suggests that the maximum value of T_a occurs when $L \approx M + 2M_f$. Figure 4 shows that same qualitative behaviour also persists in the presence of ferromagnetic leads.

In conclusion, we have shown that in contrast with the tunneling regime, where T_a is exponentially small, the non-local current can be dramatically enhanced by including a normal region between the F leads and superconductor. In this case the non-local current no longer decays monotonically with M or L , but instead exhibits a maximum corresponding to a square normal region with $L \approx M + 2M_f$.

Acknowledgements

Work supported in part by the European Community's Human Potential Programme under contract HPRN-CT-2000-00144, Nanoscale Dynamics.




Figure 1: *The hybrid nanostucture analysed.*

References

1. S. K. Upadhyay, A. Palanisami, R. N. Louie and R. A. Buhrman, Phys. Rev. Lett. **81**, 3247 (1998)
2. S. K. Upadhyay, R. N. Louie and R. A. Buhrman, Appl. Phys. Lett. **74**, 3881 (1999)
3. R. J. Soulen *et al.*, Science **282**, 85 (1998)
4. C. Fierz, S.-F. Lee, J. Bass, W. P. Pratt Jr and P. A. Schroeder, J. Phys.: Condens. Matter **2**, 9701 (1990)
5. M. D. Lawrence and N. Giordano, J. Phys.: Condens. Matter **8**, L563 (1996)
6. V. A. Vasko, V. A. Larkin *et al.*, Phys. Rev. Lett. **78**, 1134 (1997)




Figure 2: For normal leads (ie when $\epsilon_M = 0$) this figure shows Andreev transmission against length L for different contact separations M .

7. M. Giroud, H. Courtois, K. Hasselbach, D. Mailly and B. Pannetier, Phys. Rev. **B58**, R11872 (1998)
8. V. T. Petrashov, I. A. Sosnin, I. Cox, A. Parsons and C. Troadec, Phys. Rev. Lett. **83**, 3281 (1999)
9. M. D. Lawrence and N. Giordano, J. Phys.: Condens. Matter **11**, 1089 (1999)
10. F. J. Jedema, B. J. van Wees *et al.*, Phys. Rev. **B60** 16549 (1999)
11. O. Bourgeois, P. Gandit, J. Lesueur, R. Mélin, A. Sulpice, X. Grison and J. Chaussy, cond-mat/9901045
12. F. Taddei, S. Sanvito, C.J. Lambert and J.H. Jefferson, Phys. Rev. Lett. **82** 4938 (1999)
13. G. Falci, D. Feinberg and F. Hekking, Europhysics Letters **54** 255 (2001)
14. G. Deutscher and D. Feinberg. App. Phys. Lett. **76** 487 (2000)
15. G.B. Lesovik, T. Martin and G. Blatter., cond-mat/0009193
16. M. S. Choi, C. Bruder and D. Loss, Phys. Rev. **B62** 13569 (2000)
17. R. Mélin, J. Phys. Condens. Matter, **13** 6445 (2001)
18. Lambert C.J., J. Phys.: Condens. Matter, **3** 6579 (1991)
19. C.J. Lambert and R. Raimondi, J. Phys. Condensed Matter, **10** 901 (1998)
20. N.K. Allsopp, V.C. Hui, C.J. Lambert and S.J. Robinson, J. Phys.: Condens. Matter **6** 10475 (1994)




Figure 3: For normal leads, this figure shows Andreev transmission against contact separation M for different lengths L .




Figure 4: For ferromagnetic leads (ie $\epsilon_M = 3.8$) with antialigned magnetizations, this figure shows Andreev transmission length L for different contact separations M .




Figure 5: For ferromagnetic leads (ie $\epsilon_M = 3.8$) with anit-aligned magnetizations, this figure shows Andreev transmission against contact separation M for different lengths L .